

# Cambet Tablet 20mg

(Piroxicam  $\beta$  Cyclodextrin)

کیمیٹیٹ ٹیبلٹ  
۲۰ ملی گرام  
(پائی رومی کیٹیم پیٹا سائیکلو ڈیکسٹریں)

## **COMPOSITION:**

Each tablet contains:  
Piroxicam (as Beta Cyclodextrin).....20mg  
**(Innovator's Specifications)**

## **DESCRIPTION:**

It is non steroidal anti inflammatory agent with analgesic and antipyretic activity. Cambet is a new formulation of Piroxicam as a complex with cyclodextrin in the molar ratio 1:2.5. cyclodextrin, produced by enzymatic hydrolysis of common starch. Cambet is very soluble in water and has a more rapid and complete absorption than plain Piroxicam after oral or rectal administration.

## **INDICATIONS:**

Cambet is indicated for symptomatic relief of rheumatoid arthritis, osteoarthritis, ankylosing spondylitis.

## **DOSAGE & ADMINISTRATION:**

1 tablet (equivalent to 20 mg of Piroxicam) per day.

In elderly patients it may be necessary to reduce the dosage (half a tablet) and limit the duration of treatment.

## **PRECAUTIONS:**

The product must be used under strict medical control in patients with a medical history of disturbances in the upper gastrointestinal tract. Particular caution must be taken in subjects with cardiocirculatory insufficiency, arterial hypertension, reduced hepatic or renal function, previous or current blood alterations, bronchial asthma and elderly patients. Piroxicam may effect concentration and it is therefore not advisable to drive or undertake any activity requiring quick reflex action. As with other drugs having similar activity, Piroxicam may increase BUN ( Blood Urea Nitrogen) in some patients; however, BUN does not keep on increasing as the therapy continues, but reaches a steady level which goes back to or towards the norm on discontinuing the treatment. The increase of BUN is not associated with an increase of serum creatinine. Piroxicam, like other NSAIDs, decreases platelet aggregation and prolongs bleeding; this should be remembered when hematological tests are carried out and when patients undergo concomitant treatment with drugs that inhibit platelet aggregation.

## **SIDE EFFECTS:**

The most commonly found side effects are gastrointestinal disturbances which are represented by nausea, epigastric distress,

constipation and diarrhoea. Other noted side effects: signs of hypersensitivity (skin rash), headache, vertigo, asthenia, changes in blood chemistry, increase of BUN. Rare side effects: gastric ulcers with or without hemorrhages, vomiting, allergic oedema of the face and hands, increase of cutaneous photosensitivity, ocular disturbances, aplastic anemia, pancytopenia, thrombocytopenia, increase of liver function parameters, jaundice, acute renal insufficiency, water retention that may occur in the form of oedema (mainly ankle oedema), or cardiocirculatory disturbances (hypertension, congestive heart failure). Sporadic cases of gastric ulcer with perforation, Stevens-Johnson syndrome, Lyell's disease, agranulocytosis, bladder disorders, shock and warning symptoms, acute heart failure, stomatitis, alopecia and nail growth disorders have been associated with the use of Piroxicam.

## **CONTRAINDICATIONS:**

Piroxicam must not be used in subjects known to be hypersensitive to the drug, nor in subjects with gastroduodenal ulcer, gastritis, dyspepsia, severe hepatic or renal disturbances, severe heart failure, severe hypertension, severe blood alterations or hemorrhagic diathesis. It is possible that cross sensitivity with acetylsalicylic acid or other NSAIDs exist. There fore, Piroxicam must not be administered to patients in whom acetylsalicylic acid or other NSAIDs induce the symptoms of asthma, rhinitis or urticaria. The product is contraindicated in ascertained or suspected pregnancy, during lactation and in children.

## **OVERDOSAGE:**

In the event of any overdosage with Cambet a supportive and symptomatic therapy should be given.

## **STORAGE & INSTRUCTIONS:**

Store between 15-25°C.

Protect from heat, sunlight & moisture.

Keep away from the reach of children.

**To be sold on the prescription of a registered medical practitioner only.**

## **HOW SUPPLIED:**

Cambet 20mg Tablet  
20's Tablets.

خوراک و طریقہ استعمال:

ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات:

دو دوکے ۲۵ ڈاکٹر کی ہدایت پر پورے درجہ حرارت کے درمیان رکھیں۔

صوبہ گری اور مٹی سے بچیں۔ بچوں کی پہنچ سے دور رکھیں۔

صرف رجسٹرڈ ڈاکٹر کے نسخے کے مطابق فروخت کریں۔

Manufactured by:  
**PHARMASOL**  
PRIVATE LIMITED  
Plot # 549, Sundar Industrial Estate,  
Lahore, Pakistan.