

120mm

Flouramet Sterile Ophthalmic Suspension

(Fluorometholone)

فلورامیٹ
جرائیم سے پاک
آنکھوں کے قطرے
(فلورومیتھولون)

COMPOSITION:

Flouramet Sterile Ophthalmic Suspension 1mg

Each ml contains:

Fluorometholone1mg

(USP Specifications)

Flouramet Sterile Ophthalmic Suspension 2.5mg

Each ml contains:

Fluorometholone2.5mg

(USP Specifications)

DESCRIPTION:

FLOURAMET consists of Fluorometholone as an active ingredient. It is a glucocorticoid employed, usually as eye drops. Chemically fluorometholone is 9-Fluoro-11β, 17-dihydroxy-6α-methylpregna-1, 4-diene-3, 20-Dione. Fluorometholone ophthalmic suspension is indicated for use in the treatment of steroid responsive inflammatory conditions of the palpebral and bulbar conjunctiva, cornea, and anterior segment of the eye.

MECHANISM OF ACTION:

Inhibition of the inflammatory response to inciting agents of mechanical, chemical or immunological nature. Generally accepted explanation of this steroid property has not been advanced. However, corticosteroids are thought to act by the induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A2. Adreno corticosteroids and their derivatives are capable of producing a rise in intraocular pressure. In clinical studies on patient's eyes treated with both dexamethasone and fluorometholone, Fluorometholone demonstrated a lower propensity to increase intraocular pressure than did dexamethasone.

INDICATIONS:

Flouramet is indicated in steroid responsive inflammation of the palpebral and bulbar conjunctiva, cornea and anterior segment of the globe.

They are also indicated for the treatment of allergic and vernal conjunctivitis

DOSAGE AND ADMINISTRATION:

1 to 2 drops instilled into the conjunctival sac two to four times daily. During the initial 24 to 48 hours the dosage may be safely increased to 2 drops every hour. Care should be taken not to discontinue therapy prematurely.

PRECAUTIONS:

The use of a corticosteroid medication in the treatment of patients with a history of herpes simplex keratitis requires great caution. Prolonged use may cause increased intraocular pressure in susceptible individuals resulting in glaucoma, with damage to the optic nerve, defects in visual acuity and fields of vision, posterior sub-capsular cataract formation, or may aid in the establishment of secondary ocular infections from fungi or viruses liberated from ocular tissues. Various ocular diseases and long term use of topical corticosteroids has been known to cause corneal and scleral thinning. Use of topical corticosteroids in the presence of thin corneal or scleral tissue may lead to perforation. Acute purulent untreated infection of the eye may be masked or activity enhanced by presence of steroid medication. Safety and

effectiveness have not been demonstrated in children of the age group 2 years or more.

As fungal infections of the cornea are particularly prone to develop, coincidentally with long term local steroid applications, fungus invasion must be suspected in any persistent corneal ulceration where a steroid has been used or is in use. Intra ocular pressure should be checked frequently.

PREGNANCY & LACTATION:

There are no adequate and well controlled studies available. The drug should be used during pregnancy and lactation only if the potential benefit justifies the anticipated risk or as suggested by the treating physician.

SIDE EFFECTS:

Elevation of intraocular pressure (IOP) with possible development of glaucoma, and infrequent optic nerve damage, posterior subcapsular cataract formation, secondary ocular infection from pathogens liberated from ocular tissues, perforation of the globe and delayed wound healing.

CONTRAINDICATIONS:

- Acute superficial herpes simplex keratitis.
- Fungal diseases of ocular structures.
- Vaccinia, Varicella and most other viral diseases of the cornea and conjunctiva.
- Tuberculosis of the eye.
- Hypersensitivity to any of the constituents of the medication.

STORAGE & INSTRUCTIONS:

Store between 15-25°C. Protect from heat, sunlight, moisture and do not freeze. Keep away from the reach of children. Use within one month after first opening the bottle and discard the remaining portion.

Do not touch the dropper tip to any surface as this may contaminate the suspension.

To be sold on the prescription of a registered medical practitioner only.

SHAKE WELL BEFORE USE.

For ophthalmic use only.

HOW SUPPLIED:

Flouramet Sterile Ophthalmic Suspension 1mg

5ml sterile ophthalmic suspension in a plastic dropper bottle.

Flouramet Sterile Ophthalmic Suspension 2.5mg

5ml sterile ophthalmic suspension in a plastic dropper bottle.

خوراک و ہدایات:

ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

دوا کو جگہ دہری ہتھی کریدو جب حرارت کے درمیان رکھیں۔

جب آپ آرکی ٹی اور جگہ ہونے سے بچائیں۔ بچان کی حفاظت سے رکھیں۔

کبھی دھو ہاتھ نہ لیں۔ دوا کو ایک جگہ استعمال کریں۔

استعمال سے پہلے ہاتھ جھینس کر لیں۔

صرف مستعدہ آنکھ کے لئے کے مطابق فرودت کریں۔

Manufactured by:

PHARMASOL

PRIVATE LIMITED

Plot # 549, Sundar Industrial Estate, Lahore, Pakistan.

PM-0580-00

170mm

* Sign. Client

* Sign. Marketing Deptt. Shannza

* Sign. Designer Shannza

* for Design, Text, Size, Color & Color Placement

* Sign. Head of Design Deptt. Shannza